

SCIENTIFIC RESEARCH (INDEX)

JOURNAL OF GERONTOLOGY AND GERIATRIC MEDICINE, 2021

MMRC (Alhydran), an interesting treatment option for a disrupted skin barrier	13
---	----

BURNS 2017

Moisturisers in scar management following burn: A survey report	14
---	----

JOURNAL OF GERONTOLOGY AND GERIATRIC RESEARCH, 2017

Application of medical moisture retention cream (ALHYDRAN), a new option in the treatment of Venous Eczema	15
--	----

BURNS 2013

Application of medical moisture retention cream (ALHYDRAN), a new option in the treatment of Venous Eczema	16
--	----

CLINICAL STUDIES (INDEX)

EBA 2023

The occurrence of itch after burns: is there a need for oral medication?	17
--	----

EBA 2018

The effect of a topical cream application on water distribution in healthy skin and burn scars: pilot project	18
---	----

WUWHS 2016

ALHYDRAN, a new option in the treatment of hypostatic eczema	19
--	----

EBA 2015

Use of ALHYDRAN in the treatment of burn scars following a chemical explosion - a case study	20
--	----

EWMA 2014

Psychological wellbeing – scar	21
--------------------------------	----

EBA 2013

Development of a new brochure making aftercare of burns more understandable for patients	22
--	----

EBA 2011

Application of Alhydran in the treatment of pathologic scars after cryoshape therapy	23
Scars after large surface thermal burns-registration of quality of life and therapeutic influence of Alhydran	24

DAV 2010

Effect of scar treatment products on unpleasant physical manifestations in patients suffering from extensive burn scarring	25
--	----

EBA 2007 / ISBI 2008

Itching, is there a remedy? A pilot study with a rehydration gel-crème in burn patients	26
---	----

EBA 2007

Two years of experience with a new product based on aloe vera for hydration of burn scars	27
---	----



van Leen MWF, et al. J Gerontol Geriatr Med 2021, 7: 083

DOI: 10.24966/GGM-8662/100083

HSOA Journal of Gerontology and Geriatric Medicine

Research Article

MMRC (Alhydran®), an Interesting Treatment Option for a Disrupted Skin Barrier

van Leen MWF^{1*}, Hoeksema H^{2,3} and Schols JMGA⁴¹Elderly care physician, Rotterdam, The Netherlands²Burn Center, Ghent University Hospital, Gent, Belgium³Department of Plastic Surgery, Ghent University Hospital, Gent, Belgium⁴Caphri Health Services Research, Maastricht University, Maastricht, The Netherlands

Abstract

Skin aging, scar formation and some types of eczema are associated with dysregulation of the intracellular water content of the epidermis. One of the main reasons for this is a disruption of the skin barrier function which subsequently leads to an increased Trans Epidermal Water Loss (TEWL). This often results in a dry, fragile, and itchy skin, requiring rehydration and restoration of the barrier function. Based on the currently available literature, there is no clear evidence about the most optimal treatment to achieve these goals and a myriad of indifferent creams are advised in (inter) national guidelines. Since 2005 a Medical Moisture Retention Cream (MMRC) Alhydran® is used in many health care settings worldwide for patients with various pathologies associated with an impaired barrier function of the skin. This article reports specifically about the available literature on MMRC and explains how its use may result in less itching and a better hydration status of the skin.

Keywords: Disrupted skin barrier; Dry skin; Indifferent cream; Itching; Loss; Natural moisturizing factor


Conclusion

An impaired barrier function of the skin occurs in different pathologies. ALHYDRAN is a moisturizer which increases skin hydration and reduces abundant TEWL. This may contribute to:

- ✓ Preservation of an adequate skin barrier function
- ✓ Improvement of Quality of Life (in patients with a disturbed skin barrier function in different pathologies).

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/burns

Moisturisers in scar management following burn: A survey report



Tanja Klotz ^{a,*}, Rochelle Kurmis ^a, Zachary Munn ^b, Kathryn Heath ^a,
John Greenwood ^a

^a Adult Burn Service, Royal Adelaide Hospital, Adelaide, South Australia, Australia

^b Joanna Briggs Institute, Faculty of Health Sciences, University of Adelaide, Adelaide, South Australia, Australia

ARTICLE INFO

Article history:

Accepted 9 January 2017

Keywords:

Burns
Emollient
Massage
Moisturiser
Transepidermal water loss

ABSTRACT

Scar management is a recognised key component of rehabilitation following burn. Moisturising often combined with massage is commenced once healing tissue has gained sufficient strength to tolerate surface friction, with the aim being to hydrate the dry scar. The studies on various moisturisers and creams provide some guidance on moisturiser selection, but many are inconclusive.

Objective: This survey aimed to determine the current expert opinion regarding moisturiser recommendations, including the basis for these recommendations, across the burns community.

Methods: A brief web-based survey was distributed to burn therapists via mailing lists of the Australian and New Zealand Burn Association (ANZBA), and American Burn Association (ABA) 'Occupational and Physical Therapist Burn Special Interest Group'.

Results: The fifty three respondents indicated that there were 29 different moisturisers commonly recommended in practice. Three main themes were indicated as influencing recommendations for moisturiser: the perceived effects on the scar/skin (48%); the general properties of the moisturiser (38%); the ingredients (14%). Therapists reported that the principle stimuli determining their recommendations were patient feedback and the choice of the previous burn therapist in their service. Many were also guided by medical staff, pharmacists and sales representatives. Only three respondents were able to provide citations for published evidence supporting their recommendations.

Conclusions: There is a paucity of evidence currently to support optimal moisturiser choice. This survey demonstrates that conflicting opinions are held on the ideal moisturiser brand, properties and ingredients. The recommendations made are based on low level evidence. Further research is required to inform clinicians which moisturiser to recommend to their clients. An ideal moisturiser should be one that is conducive to scar maturation, non- or minimally irritant, prevent skin drying, minimise transepidermal water loss and have no negative effect on barrier function.

Conclusion

- ✓ TEWL appears to be one of the most important considerations regarding future research in the area of moisturisers effect on scars
- ✓ The ideal moisturizer should be conducive to scar maturation, be low irritant, prevent skin drying and negative effects to barrier function, and minimize TEWL
- ✓ Avoid watery creams. Aqueous creams increases TEWL
- ✓ ALHYDRAN has demonstrated to have effect on TEWL and potentially the scar outcome
- ✓ More studies of similar quality and content such as that assessing ALHYDRAN are required



Journal of Gerontology & Geriatric Research

Rondas and Schols, J Gerontol Geriatr Res 2017,
6:1
DOI: 10.4172/2167-7182.1000395

Research Article

OMICS International

Application of Medical Moisture Retention Cream (ALHYDRAN®), A New Option in the Treatment of Venous Eczema

Rondas AALM^{1,2*} and Schols JMGA¹

¹Maastricht University, Department of HSR - CAPHRI, Care and Public Health Research Institute, Maastricht, The Netherlands

²Kenniscentrum Wondbehandeling, De Zorggroep, Venray, The Netherlands

*Corresponding author: Rondas AALM, Maastricht University, Department of HSR - CAPHRI, Care and Public Health Research Institute, Maastricht, Kentstraat 20, 6137 JT Sittard, The Netherlands, Tel: +31478514044; E-mail: a.rondas@maastrichtuniversity.nl

Rec date: Jan 23, 2017; Acc date: Feb 07, 2017; Pub date: Feb 09, 2017

Copyright: © 2017 Rondas AALM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Aim: Patients with venous eczema may suffer considerably from redness, crusts, pain, flaking and itching. In general, as treatment, compression therapy and indifferent ointments/crèmes are used, often together with topical steroids, though the latter may exhibit considerable side effects. This study aims to explore the effect of Medical Moisture Retention Cream (MMRC = ALHYDRAN®) on the symptoms and complaints of patients suffering a venous eczema, often next to an existing VLU. The working mechanism of MMRC involves a combination of the moisturizing effect of Aloe Vera gel and the moderate occlusion effect of added fatty acids.

Method: In an open case series, 18 patients attending an outpatient wound clinic with moderate venous eczema (maximum TIS score < 3) participated. MMRC was applied twice a day and its use was clinically assessed for 4 consecutive weeks. Next to the registration of patient characteristics and capturing the clinical details of the skin lesions in weekly pictures, a VAS scale to assess the patients' and caregivers' experience with the treatment was used.

Results: During the application of MMRC, the signs and symptoms of a dry, itchy, scaly, crusty and erythematous skin, faded in all patients. The skin condition of most participating patients improved and there was also a visibly better skin hydration status in all patients. Relevant aspects such as 'night rest', 'mood' and 'social participation' improved, except for 2 patients. The clinical skin condition of one of these deteriorated in a week, which was not related to the use of MMRC. No side effects to the use of MMRC were observed. The wound care experienced nurses assessed MMRC as effective and feasible.


Conclusion: This study shows that MMRC is effective, safe, and feasible in the treatment of venous eczema. Future Randomized Controlled Trials are necessary to compare the efficacy and feasibility of MMRC with the application of other hydrating creams/ointments and topical steroids.

Conclusion

- ✓ This study shows that ALHYDRAN is effective, safe and feasible in the treatment of venous eczema
- ✓ During the application of ALHYDRAN, the signs and symptoms of a dry, itchy, scaly, crusty and erythematous skin, diminished in all patients
- ✓ Visibly improved skin hydration status in all patients
- ✓ The skin condition of most participating patients improved
- ✓ Relevant aspects such as 'night rest, mood and social participation' improved in most participants

Available online at www.sciencedirect.com

SciVerse ScienceDirect

journal homepage: www.elsevier.com/locate/burns

Scar management by means of occlusion and hydration: A comparative study of silicones versus a hydrating gel-cream

Henk Hoeksema^{1,*}, Marie De Vos¹, Jozef Verbelen, Ali Pirayesh, Stan Monstrey

Department of Plastic and Reconstructive Surgery – Burn Centre, Ghent University Hospital, Ghent, Belgium

ARTICLE INFO

Article history:

Accepted 29 March 2013

Keywords:

Scar
Scar treatment
Silicone
Silicone gel
Fluid silicone gel
Silicone gel sheets
Hydrating gel-cream
Moisturizers
Hydration
Occlusion
Trans epidermal water loss

ABSTRACT

Despite the worldwide use of silicones in scar management, its exact working mechanism based on a balanced occlusion and hydration, is still not completely elucidated. Moreover, it seems peculiar that silicones with completely different occlusive and hydrating properties still could provide a similar therapeutic effect.

The objective of the first part of this study was to compare the occlusive and hydrating properties of three fluid silicone gels and a hydrating gel-cream. In a second part of the study these results were compared with those of silicone gel sheets.


Tape stripped skin was used as a standardized scar like model on both forearms of 40 healthy volunteers. At specific times, trans epidermal water loss (TEWL) and the hydration state of the stratum corneum were measured and compared with intact skin and a scar-like control over a 3–4 h period.

Our study clearly demonstrated that fluid silicone gels and a hydrating gel-cream have comparable occlusive and hydrating properties while silicone gel sheets are much more occlusive, reducing TEWL values far below those of normal skin.

A well-balanced, hydrating gel-cream can provide the same occlusive and hydrating properties as fluid silicone gels, suggesting that it could eventually replace silicones in scar treatment.

© 2013 Elsevier Ltd and ISBI. All rights reserved.


ELSEVIER



Conclusion

ALHYDRAN: A well-balanced, hydrating cream with occlusive and hydrating properties for treatment of disrupted skin barrier (e.g. scars).

- ✓ Significantly reduced TEWL
- ✓ Increased stratum corneum water content
- ✓ Long-lasting effect



Burns 2013:
test areas

The occurrence of itch after burns: is there a need for oral medication?



H.W.C. Hofland RN, SCRN, MScN Burn center Maasstadziekenhuis, Rotterdam, the Netherlands.



Correspondence:
hoflandh@maasstadziekenhuis.nl

Introduction:

This study describes the occurrence and management of itch in adult patients admitted to the Burn Center in Rotterdam, with a follow up for one year.

Methods:

For one month all adult patients admitted were asked to participate in the study. After the first episode of itching the Burn Itch Questionnaire (BIQ) was used to record itch frequency until discharge. From the patient's medical dossier data were extracted on medical history, prescribed medication and routine usual care. Follow up was for 3, 6 and 12 months after discharge. The itch was scored with the Visual Analogue Thermometer (VAT); 0 no itch - 10 worst itch.

Results N=35


Male 80%
Mean LOS 21 days (3-78)
Follow up 86%
Mean age 42 yr (18-81)
Flame burns 22 (63%)
Surgery 15 (43%)
Mean TBSA 9% (1-31)

First itch (3-25 days)

VAT 0-1 7 (20%)
VAT 2-5 24 (69%)
VAT 6-8 4 (11%)
VAT 9-10 0 (0%)

After three and six months, 32% and 40% of the respondents were without pruritus, respectively; after 12 months this was 60%. After 12 months 34% still applied this ointment which was preferred to oral medication. After 12 months, only one patient used anti-histamines.


Use of medication (%)



Effect of medication (%)

3 months (26%) no improvement (33%)
moderate effect (56%)
complete reduction (11%)
6 months (12%) moderate effect
12 months (3%) moderate effect


Use of Alhydran (%)



Effect of Alhydran (%)

3 months (80%) no improvement (7%)
moderate effect (18%)
complete reduction (75%)
6 months (49%) no improvement (17%)
moderate effect (11%)
complete reduction (72%)
12 months (34%) no improvement (9%)
moderate effect (18%)
complete reduction (73%)

Itch complaints (%)



Conclusion:

Pruritus is common after burn injury. 29% of the respondents still experience itch after one year; oral medication is often not needed; Topical ointments such as Alhydran are preferred.



Usual care of itch

- Itch assessment - VAT
- Medication - antihistamines (promethazine)
- Healed skin - application of Alhydran



Alhydran

- Medical cream based on Aloe Vera
- Moisture-regulating cream
- Easy application
- Reduces dry and reddened skin
- Rapidly reduces itching
- Increases skin elasticity

Factors increasing or inducing itch

- Dry skin
- Too warm (sweating, weather, cloths)
- Stress and emotions
- Standing in upright position
- Removal pressure garments
- Itch comes with spells
- Does sometimes interfere with daily activities and sleep

CLINICAL STUDY

Presented at the European Burn Association (EBA), Rotterdam, the Netherlands, 2018

OSCARE - Organisation for burns, scar after-care and research, Antwerp, Belgium

The effect of a topical cream application on water distribution in healthy skin and burn scars: pilot project

Authors: P. Moortgat ¹, M. Anthonissen ^{1,2}, J. Meirte ^{1,3}, U. Van Daele ^{1,3}, T. Vanhullebusch ^{1,3}, C. Lafaire ^{1,4}, L. De Cuyper ^{1,4}, K. Maertens ^{1,5}



- ¹ OSCARE, Organisation for burns, scar after-care and research, Antwerp, Belgium
² KU Leuven, Department of Rehabilitation Sciences, Belgium

- ³ University of Antwerp, Rehabilitation Sciences and Physiotherapy, Belgium
⁴ ZNA Stuivenberg, Burns Center, Antwerp, Belgium
⁵ Vrije Universiteit Brussel, Department of Clinical and Lifespan Psychology, Belgium

Objectives:

Moisturizing is an essential part of scar management. However, the evidence for topical applications is poor or lacking in skin and scar research. The study aimed to investigate baseline values of water distribution in healthy and scarred skin and to examine the effect of Alhydram® (fig. 1) on water distribution in healthy and scarred skin over time.



Fig. 1: Alhydram® BAP Medical B.V.



Fig. 2: Corneometer® probe MPA 580 Courage Khazaka

Methods:

For this pilot project, 20 healthy skin sites and 20 burn scar sites were included. At baseline, all sites were assessed for water content, using the Corneometer® (fig. 2) ¹ and for water distribution, using the Moisture Map® (fig. 3) ¹. Water distribution is represented using gray index (derived from original image), corner density, number of intersection lines and number of cells (calculated from segmented image) (fig. 4).



Fig. 3: Moisture Map® Courage Khazaka




Fig. 4: Original image (left) is transferred into a segmented image (right)

Half of the healthy skin sites and scar sites formed the intervention group and were hydrated with Alhydram®, while the other half served as controls. Thirty minutes after application, these sites were re-assessed similarly. For the statistical analysis, an independent samples T-test was carried out for the between-group analysis of baseline values. A paired sample T-test was used for the within-group analyses over time.

References:

- 1) www.courage-khazaka.de
 2) www.alhydram.com/professionals-doctors-nurses-paramedics/

Results:

The baseline water content values and water distribution parameters for the healthy skin sites and scar sites are presented in table 1. Statistically significant differences between healthy skin and scar sites are found for water content and water distribution (except for gray index) at baseline ($p \leq .001$) (table 1).

Table 1: Between-group analysis (p-value) at baseline for water content values and water distribution parameters of healthy skin sites and scar sites (mean \pm standard deviation).

	Healthy skin site	Scar site	p-value
Water content (AU)	37.3 \pm 5.6	28.3 \pm 9.9	.001*
Gray index	21.7 \pm 5.9	18.5 \pm 8.9	.195*
Corner density (%)	3.2 \pm 0.7	1.5 \pm 0.8	.000*
Intersection lines	679.2 \pm 122.8	359.3 \pm 153.8	.000*
Cells	357.0 \pm 75.6	183.9 \pm 83.5	.000*

Thirty minutes after Alhydram® application (T30), all outcome measures of healthy skin improve significantly compared to baseline (T0) ($p \leq .018$). For the scarred skin a trend towards improvement was noticed, however not significant (ns) (table 2).

Table 2: Within-group analysis over time for the healthy skin sites and scar sites (p-value) with mean difference (T30-T0) \pm standard deviation (sd).

	Healthy skin site		Scar site	
	mean T30-T0 \pm sd	p-value	mean T30-T0 \pm sd	p-value
Water content (AU)	4.1 \pm 2.6	.001*	3.5 \pm 9.3	ns
Gray index	4.4 \pm 4.8	.018*	1.8 \pm 5.4	ns
Corner density (%)	0.9 \pm 0.7	.003*	0.2 \pm 0.6	ns
Intersection lines	151.5 \pm 132.2	.006*	45.3 \pm 114.5	ns
Cells	102.6 \pm 76.6	.002*	25.7 \pm 67.2	ns

Conclusion:

The uptake of water after moisturizing differs significantly between healthy skin sites and scar sites. A single Alhydram® application shows a significant increase in water content and water distribution after 30 minutes in healthy skin.

In scarred skin, there is a trend in improvement. These results support previous findings that scarred skin is less permeable for a single application than healthy skin. For future research, we would suggest multiple Alhydram® applications (at least 3x / day, as described in usage guidelines of Alhydram®) on healthy skin sites and scar sites with a longer follow-up period ².

Presented at the 2nd European Burns Association educational course (EBA), Rotterdam, the Netherlands, 28 & 29 September, 2018

ALHYDRAN[®], a new option in the treatment of hypostatic eczema

Authors: Rondas AALM^{1,2}, Vestjens JA¹, Schols JMGA².

¹ De Zorggroep, Kenniscentrum Wondbehandeling, Venray, the Netherlands.

² Maastricht University, CAPHRI, School of Public Health and Primary Care, the Netherlands.



Maastricht University



Kenniscentrum Wondbehandeling

Introduction

Hypostatic eczema is the result of an existing chronic venous insufficiency. Patients may experience redness, crusts, pain, flaking and itching. As treatment, compression therapy and indifferent ointments and crèmes are used, often together with the application of topical corticosteroids.

The aims

In this case series the effect of a strong hydrating cream ALHYDRAN[®] (>45% Aloë Vera gel) was explored. By applying ALHYDRAN[®] it is expected that the patient's skin will become better hydrated and that further water loss is prevented. Subsequently, the topical use of corticosteroids might be less necessary.

Methods

Patients attending an outpatient wound clinic and suffering from moderate hypostatic eczema (maximum Three-Item-Severity (TIS) score <3) were included. They all used compression therapy.

ALHYDRAN[®] was applied at least twice a day and its use was clinically assessed for a maximum of 4 consecutive weeks. Next to the registration of patient characteristics and capturing the clinical details of the skin lesion in weekly pictures, a tool to assess the professionals' experience with the treatment was used.

Results

In total 18 patients, 10 men and 8 females, were included with a mean of 69.8 years. In 17 patients professionals evaluated the overall treatment effect of ALHYDRAN[®] as positive. With regard to their 'night rest', 'social participation' or 'mood', 16 patients had no complaints at the start of the application. During application the night rest and social participation of 2 patients greatly improved.

Evaluation of the effectivity and feasibility of applying ALHYDRAN[®] professionally scored respectively 6.75 and 7.5 on a scale of 10. Only 1 patient showed a much more scaly skin after one week of use and in this patient precautionally the ALHYDRAN[®] was stopped.

Conclusion

Applying ALHYDRAN[®] in patients having an itchy, scaly and dry skin because of underlying chronic venous insufficiency, may decrease the (often undesired) topical use of corticosteroids.

Start ALHYDRAN[®] treatment After four weeks of ALHYDRAN[®] treatment



Start ALHYDRAN[®] treatment After four weeks of ALHYDRAN[®] treatment



Start ALHYDRAN[®] treatment After four weeks of ALHYDRAN[®] treatment



Start ALHYDRAN[®] treatment After four weeks of ALHYDRAN[®] treatment



Start ALHYDRAN[®] treatment After four weeks of ALHYDRAN[®] treatment



Start ALHYDRAN[®] treatment After four weeks of ALHYDRAN[®] treatment



Start ALHYDRAN[®] treatment After four weeks of ALHYDRAN[®] treatment



Start ALHYDRAN[®] treatment After four weeks of ALHYDRAN[®] treatment



Presented at the World Union of Wound Healing Societies (WUWHS), Florence, Italy, September 25-29, 2016

ALHYDRAN[®] is a product of BAP Medical - Medical care for the skin. More information: WWW.ALHYDRAN.CO.UK or WWW.ALHYDRAN.NL

CLINICAL STUDY

Presented at the European Burn Association (EBA), Hannover, Germany, 2015

Scar and Burns Aftercare Department, Oosterbeek, the Netherlands

Use of ALHYDRAN cream in the treatment of burn scars following a chemical explosion - a case study

Author: Rina Rijkenberg, Bachelor of Health, Skin Therapist. Skin Therapy Clinic, Scars and Burns Aftercare Department, Oosterbeek, The Netherlands.

Background

A 58 year old male had been injured in a chemical explosion. The explosion caused a fire, resulting in flame burns to the skin on his face and hands. Burns to the face were second degree, with second to third-degree burns on the hands. On admission to hospital, the Burns Centre Maasstad in Rotterdam, treatment of the burns involved the daily use of silver sulfadiazine cream and changing the bandages until the wounds closed.

The Medical Consultant at the Burn Centre recommended continuation of the treatment by a skin therapist specialising in the aftercare of burn injuries.

Complaints

The patient complained of extremely dry and itching skin on the face and hands. His skin felt extremely tight due to the burns and his hands were highly oedematous.

Post-burn skin therapy

Our therapy consisted of twice-weekly, manual lymphatic drainage, bandaging and endermotherapy, followed by the application of silicon sheets (BAPSCARCARE). A hydrating cream (ALHYDRAN) was also applied 4 to 6 times a day as a very thin layer, by the patient himself. (Normal application is 3 times a day.) We also applied this hydrating cream before and after every endermotherapy treatment.

Results

ALHYDRAN made the patient's facial skin more flexible. The patient reported noticing a difference one or two hours after applying the hydrating cream to his face. His skin felt much looser. The following day the skin on his face was more flexible and "felt great!" His face healed completely, and the skin on his hands became more flexible. Application of the hydrating cream enabled the skin to be moved freely, without tearing, during lymphatic drainage and endermotherapy, which also contributed to the healing process. The cream also significantly reduced the itching.

After three months the skin had noticeably improved; it looked healthier and there were fewer squamae. The hydration level of the skin was improved and the patient reported less itching. Treatment was concluded after 18 months.

References

- 1) "Two years experience with a new product for the hydration of Burn Scars"; EBA Congress 2007; Monstrey, S., MD, PhD, Burns Centre, Ghent University Hospital, Belgium.
- 2) "Itching, is there a remedy?"; pilot study on rehydrating gel-cream for burns patients. EBA 2007; H.J. van Kempen, RN; Burns Centre, Rijnmond-Zuid, Rotterdam, The Netherlands.
- 3) "Scar management by means of occlusion and hydration: A comparative study of silicones versus hydrating gel-cream". Burns. 2013 Apr 29. pii: S0305-4179(13)00105-8. doi: 10.1016/j.burns.2013.03.025; Hoeksema H, Monstrey S, et al. Department of Plastic and Reconstructive Surgery - Burns Centre, Ghent University Hospital, Belgium.

Conclusions

The burns were successfully treated using a hydrating cream in combination with lymphatic drainage, bandaging and endermotherapy. The cream is pleasant to use and only a thin layer is required.

We recommend ALHYDRAN to all our burn patients, because of the excellent clinical outcomes we have achieved over the past few years with regard to itching^{1,2}, elasticity¹ and redness¹. This is due to the proven hydration³ and occlusion³ properties of this cream.



Picture 1: These pictures were taken the first time we saw the patient



Picture 2: These pictures were taken three months after commencing our treatment



Picture 3: These pictures were taken six months after commencing our treatment

De Huidtherapeuten is a skin and oedema therapy practice. Our therapists are qualified in the treatment of burns and hold the 'Skin Therapy After Burns' diploma from the University of Applied Sciences, Utrecht, which collaborates with several burns centres in the Netherlands and Belgium.*

.....
NETWORK ► HUIDTHERAPIE
.....
*NETWERKPARTNER VAN DE BRANDWONDEN STICHTING


De Huidtherapeuten
PRAKTIJK VOOR HUID - EN DEDEEMTHERAPIE

PSYCHOLOGICAL WELLBEING – SCAR

Authors: Stuart Ashman, Sara-Jane Kray, Tyne & Wear, UK

Aim

To review whether psychological wellbeing can be enhanced by improving the appearance of scar tissue, post injury.

Background

Scar management is not routinely administered for trauma wounds that occur during childhood, however, they can have an effect on how an individual perceives themselves and impact on how they dress.

Scars are the result of an injury to both the dermis and epidermis (and/or deeper tissue). Post proliferation the skin attempts to normalise during the final maturation stage of healing.

Normal remodelling of the skin can take up to 12 months, and depending on how the wound was left to heal, e.g left open or closed with sutures, can also impact on the overall outcome. Scars generally improve over time, becoming less red/pink as blood vessels are removed from the area through apoptosis and flatten as the balance between the synthesis and degradation of collagen is completed.

Routine scar management^[1] is primarily based on the use of silicones following wound closure to occlude and hydrate the scar tissue thus reducing the production of excessive collagen^[2].

Methodology

This case study reviewed the use of a silicone gel sheet and hydration gel-cream and how improvements in scar tissue can have a positive impact on a child's psychological wellbeing.

A girl aged 7, sustained a traumatic injury to her right shin, measuring 4x2cm, requiring 6 sutures (2nd August 2013, Photo 1). Following suture removal the wound dehiscd and the resulting scar area remained bright red and raised. The scars prominence resulted in the child refusing to wear dresses or shorts throughout the summer.

A dual therapy was initiated (November 2013, Photo 2) comprising a silicone gel sheet* and a hydrating gel-cream*. Tolerance to the silicone was built up to recommended wear times and the hydrating gel-cream was applied in between with an hour between the application of the hydrating gel-cream and the silicone sheet to ensure optimum results.



Photo 1 - Aug 2013



Photo 2 - Nov 2013



Photo 3 - Jan 2014



Photo 4 - Apr 2014

Results

Within 6 weeks there were considerable improvements in the elasticity, smoothness, thickness and scar appearance (Photo 3). After a total of 5 months of dual therapy the scar had significantly decreased in size and the overall appearance of the scar was flatter, paler and acceptable to the child, (Photo 4).

The child applied the products herself, stating "It makes me feel better, it's working. I put it on myself and I try to remember to do it every day. I'm really happy that the scar is fading so that I can wear nice dresses next summer without the ugly scar".

Conclusion

Scar management should be initiated immediately after wound closure and should continue until the scar no longer causes a problem both physically and psychologically. Silicones gel sheets are shown to provide pressure at the site of scarring and may be used with pressure garments. The hydration gel-cream has been shown to provide the same occlusion as silicone gels whilst also providing hydration, and improving the overall appearance of the scar. Could a combined therapy of both a silicone gel sheet and a hydration gel-cream therefore be the future?

References

1. Thomas A. Mustoe, M.D., Rodney D. Cooter, M.D., et al. (2001) International Clinical Recommendations on Scar Management, Plastic and Reconstructive Surgery, Vol 110, No. 2 (560-571)
2. Hoeksema H, et al. (2013) Scar management by means of occlusion and hydration: A comparative study of silicones versus a hydrating gel-cream. Burns. <http://dx.doi.org/10.1016/j.burns.2013.03.025>

CLINICAL STUDY

Presented at the European Burn Association (EBA), Vienna, Austria, 2013

Burn Centre, University Hospital Ghent, Belgium

Development of a new brochure making aftercare of burns more understandable for patients

Henk Hoeksema, Katrien Van Gastel, Veerle Van Geertruy, Pieter Lafaie, Jozef Verbelen, Stan Monstrey
Burn Centre, Ghent University Hospital, Belgium

Introduction:

One of the first questions of burn victims and relatives is whether they will end up with visible scars. Scars are an unavoidable consequence of dermal burns, but the outcome depends largely on the method of scar treatment. This very long and tedious process can only lead to a good result by optimal professional skills and compliance of the patient. We noticed the growing need for a patient brochure concerning the aftercare of burns because an optimal outcome can only be achieved if the patient and family completely understands the reasons behind the treatment.

Methods:




Fig 1: healing time or healing potential

		hydration	UV-protection	Pressure garments	silicones
A	HP <14 days	yes	yes	no	no
B	HP 14-18 days	yes	yes	no	no
C	HP 17-21 days	Yes	Yes	Yes	yes
D	HP >21 days	Yes	Yes	Yes	yes

Fig 2: aftercare related to healing time



Fig 3: Individual aftercare schedule



Results:

The creation of a brochure, starting with a short explanation about scar development after burns. A subdivision into three groups based on healing potential by laser Doppler imaging and exact healing time is linked to the appropriate scar treatment. With help of a figure in the brochure, showing the exact healing times for the different burned areas, it is easy for the patient to understand what kind of scar treatment is recommended. Finally answers to the most common questions are provided.

Conclusion:

This brochure improves burn patients understanding of scar treatment and improves patients compliance to the suggested treatment. As a consequence, functional and aesthetic outcomes are positively influenced.


383

Application of Alhydran in the treatment of pathologic scars after cryoshape therapy

Personal experience



Cryogun.



How cryoshape acts on scars.
 Core of the scar at -196 °C (-320.8°F) no solution of continuity of the skin.



Clinical case: Hypertrophic scarring after bilateral reduction mammoplasty.



Close up of the scars to be treated (right breast).



Close up of the scars to be treated (left breast).

Dr. Uberto Giovannini
 Service de Chirurgie Plastique Reconstructrice et Esthétique
 Hôpital Maternité de Metz
 1/5 Place sainte croix
 57045 Metz cedex1
 Tel + 33(0)3 87345166
 e-mail: ubertogiovannini@yahoo.com

Introduction

Aloe Vera has been used as a popular herbal medicine since ancient times for many conditions. In 1851 T and H Smith of Edinburgh discovered the active principle of the plant. Aloe Vera contains 75 active constituents: vitamins, enzymes, minerals, monoacids, and acid salicyliques. After a bibliographic research on EBSCO, Medline and Google with the key words: aloe Vera, topic, surgery and wound healing, we decided to introduce the topical utilization of Aloe Vera in the treatment of scar tissue. We decided to use Alhydran cream because of the characteristics of the pure and high concentration of the Aloe Vera, associated with high quality oils, Vitamins E and C. The cryoshape is a new technique for treating keloids and hypertrophic scars developed by Dr. Yaron Har-Shai. We present our experience with these new devices.

Materials and Methods

Technics

A specially designed cryo-needle was inserted into the long axis of the hypertrophic scars and keloids so as to maximize the volume of the hypertrophic scars and keloids to be frozen. After the hypertrophic scars and keloids were completely frozen, the cryoprobe defrosted and was withdrawn. After the treatment a dressing with Vaseline gauze was applied. The dressing was replaced after 48 hours by hydrocolloid. We started the application of Alhydran two weeks after these scars had been treated with cryoshape. The topical application was done twice a day for roughly four to six weeks.

Patients

From January 08 till July 09 we have treated 13 patients for a total of 21 scars. The age of the patients were between 17 and 62 years old

5 patients presented pathological scars after breast surgery, 2 patients after abdominal surgery, 2 patients after ear surgery, 1 patient after cleft lip surgery, 1 patient after sternotomy and one patient after burn injury.

Results

None of the patients experienced an intolerance or allergy to the Alhydran. All patients experienced improvement in the scars aspect and pain and itching presented before treatment had disappeared. No adverse events were reported by any of the patients.

Conclusion


Cryoshape and Alhydran showed a good synergy in the management of pathologic scars. Further, well-designed trials should be carried out to determine the effectiveness of Alhydran in improving scars outcome




Keloid of the trunk pre operative view




Follow up at six months after one session of cryosurgery and six weeks of application of Alhydran



Follow up at six weeks of application of Alhydran



Introducing the cryoshape through the scar



Intraoperative view: you can note the freezing of the scar





Scar completely frozen.



Immediate post operative



Pathologic scars of mediastinum sequel of cardiac surgery: pre operative view before cryosurgery



CLINICAL STUDY

Presented at the European Burn Association (EBA), The Hague, the Netherlands, 2011
Burn Rehabilitation Centre, Bad Klosterlausnitz, Germany



Moritz Klinik Bad Klosterlausnitz
Orthopädisch-traumatologische Abteilung
Reha-Zentrum für Brandverletzte

Scars after large surface thermal burns – Registration of Quality of Life and therapeutic influence of Alhydran®

Hans Ziegenthaler


Question:

Scars, as a result of large surface thermal burns, show in comparison with normal and sound skin, serious functional deficits. These deficits can be described by means of objective, biomechanical and physiological characteristics and by the subjective observation of itching, skin tension, dryness and redness of the skin. The main goal of the study was to investigate the influence of the scarring on the Quality of Life related to the patients circumstances, and to measure this as well as the influence of skin care products in the early stage of scar maturation during the rehabilitation of the patients.

Material and Method

Material:	Patient population
number (n)	75
men / woman	54 / 21
average age (years)	51,3 ± 14,6
BMI (kg/m ²)	23,9 ± 5,7
average total burn surface area (%)	23,0 ± 15,0
range accident to start inpatient rehabilitation (days)	55,1 ± 30,2
duration of treatment (days)	34,9 ± 16,1

Negative Bodyexperiences related to the burn injuries were noted in the beginning of the test in the following percentages of all patients (%)




The Pattonnaire® questionnaire is a best-suited patient questionnaire for the interactive anamnesis and allows the objective measurement of symptoms, disabilities and well-being. The patient completes the Pattonnaire questionnaire himself. It is then discussed with the physician, therapist, study attendant, supplemented and, if necessary, corrected accordingly. The Construct Validity revealed a very high agreement between the answers to the questionnaire and the personal interviews (94% of the specimen). There was also a partial conformity (6% of the specimen) by older patients with cognitive problems. The reliability by repeated completion of the questionnaire (Test-Retest) produced good values. The Intraclass Correlation Coefficients (ICC) were over 0.7 (good) by 22 of the 33 questions. The Pattonnaire measures more complaints (up to 10 symptoms) than other questionnaires (e.g. SF 36), also specific items of burn victims with a specific module. He is not only of simple design, but also clearly defined and patient-friendly.

The SF-36v2® Health Survey asks 36 questions to measure functional health and well-being from the patient's point of view. It is a practical, reliable, and valid measure of physical and mental health that can be completed in five to ten minutes. It's called a generic health survey because it can be used across age (18 and older), disease, and treatment group, as opposed to a disease-specific health survey which focuses on a particular condition or disease. The survey is meaningful to patients, clinicians, researchers, and administrators across the health care spectrum, and has various applications. The SF-36 provides scores for each of the eight health domains [Physical Functioning (PF), Role-Physical (RP), Bodily Pain (BP), General Health (GH), Vitality (VT), Social Functioning (SF), Role-Emotional (RE), Mental Health (MH)] and psychometrically-based physical component summary (PCS) and mental component summary (MCS) scores.


Data evaluation: The specifications have been analyzed by the program for statistical evaluation WINDOWS SPSS 13.0.

Results

The Pattonnaire® showed, especially in the fields of redness (20,4%) and skin tension (16,9%) a substantial improvement compared to the initial values. Also the reduction of itching (13,5%) and dryness of the skin (13,8%), were clearly noted in the scoring of the patients. The general score on the experienced impediments of the burn scars was reduced in 2/3 of the patients, and the general quality of life experience was increased with 12,6%. The use of anti-histamines in the study group of patients was reduced during the total treatment period by 1/3. Side effects of the treatment with Alhydran® were not at all experienced. (left)



As could be expected, the SF-36 short-form health survey, showed for all patients and all items and in all subscales, an underscoring of the average Norm value of 50 of the standard population. The biggest underscoring in combination to this standard normation value of 50 of the SF-36 Physical Component Summary and Mental Component Summary, were found in the scales of Physical Functioning, Role-Physical and General Health. This could be expected because of the high physical functional deficits of the patients. The physical component summary (PCS) had originally an average of 29,7 ± 6,3 which increased after 4 weeks to 21,9 ± 5,9. In the mental component summary (MCS) the starting value was 49,3 ± 7,9 and this increased to 51,7 ± 8,0. The Physical Functioning (PF) had in the beginning a value of 21,7 ± 9,5 and this improved to 21,9 ± 7,4. The biggest improvement in the patient situation was obtained with the Role-Emotional scoring (RE), where the value increased from 19,1 ± 5,9 towards 33,8 ± 7,9. The values for burn patients however still remain below the values for healthy comparative groups.(right)



Conclusions:

1. This first use of the Pattonnaire® questionnaire with a new skin module for the follow-up of burns, showed in practice stable and reliable characteristics of use in the monitoring of burn patients.
2. Alhydran® was successfully used during the early maturation of the scars, to reduce the subjective and unpleasant side effects of burn scars such as redness and skin tension, without noticeable side effects.
3. Alhydran® could trouble free be combined with (textile garments) pressure therapy. Inhomogeneous treatment groups and the missing randomization in this study, do not allow to draw conclusions beyond the observations of this study about a trend in treatment effects.
4. Further clinical, probably multi-centre studies with a larger randomized group of patients, as well as a post-study check, are needed to be able to draw statistically relevant conclusions.

Die Untersuchung fand mit freundlicher Unterstützung der Fa. BAP Medical BV statt.

Dr. med. H. Ziegenthaler Moritz Klinik GmbH & Co.KG Hermann-Sachse-Str. 46 D-07639 Bad Klosterlausnitz phone: +49 36601 49 463 Email: hans.ziegenthaler@moritz-klinik.de



28th Annual Conference
of the German-speaking Working Group
for Burn Treatment

dav2010

13–16 January 2010
Schladming – Rohrmoos (Styria – Austria)

ABSTRACT BOOK

p47

Effect of scar treatment products on unpleasant physical manifestations in patients suffering from extensive burn scarring

Lead author: Dr. med. Hans Ziegenthaler
Moritz Clinic, Burns Rehabilitation centre, Hermann-Sachse-Str. 46, 07639 Bad Klosterlausnitz

Purpose:

Scars lack endogenous sebaceous glands, which means that they do not have a physiological fatty coating to protect the tissue against acid. Consequently, the scarring process is accompanied by unpleasant physical manifestations such as itching, tightness, dryness and flaking, the subjective perception of which varies from one individual to another. The aim of our investigation is to ascertain how far the effect of scar treatment products in the early stages of scar formation during rehabilitation can be measured.

Patients and methods:

15 patients with burn injuries taking part in an in-patient rehabilitation programme (five to eight weeks after their accident) were observed as part of a clinical treatment observation study. The patients all had healed scar tissue patches on the trunk and limbs which flaked quite severely, felt tight and/or itched both at rest and while the patient was moving. Wound care was generally given every day, in some cases more than once a day, using the proven product SR Unguentum cordes. After inflammatory factors had been ruled out, those patients who were continuing to experience itching and a sensation of tightness exceeding 50% on an intensity scale were switched to ALHYDRAN® cream. The state of the scars was assessed using the Vancouver Scar Scale at point T0, T1 (14 days later) and T2 (a further 14 days later). At these points photographs were taken, medication requirements (analgesics, neuroleptics, antihistamines) were noted, patients' pain perception according to the VAS scale was recorded and their health-related quality of life (SF 36) was ascertained. The Pationnaire® questionnaire, which had never previously been used with burns patients, was also employed (this questionnaire investigates various everyday and injury-specific items). The trial group was too small for statistical assessment, so the findings were assessed by means of descriptions and photographs.

Findings:

A marked improvement compared to the baseline was observed in the subjective categories assessed, particularly itching and a sensation of tightness. However, the treatment group, the period investigated and the absence of randomisation make it impossible to reach any more far-reaching conclusions than an observation of change. No adverse effects were reported. On the other hand, convincing application properties in the burns patient group were indicated by the Pationnaire® questionnaire, which had previously been used only in the context of injuries to the supporting and locomotor systems but which now has a skin module.

Conclusions:

ALHYDRAN® appears to be able to minimise manifestations such as itching and a sensation of tightness, which patients find very distressing, in the early stages of scar formation. This form of scar treatment is well tolerated and can easily be combined with the administration of textile-based compression therapy. We intend to conduct a clinical study with a larger randomised group and a follow-up after three months in order to obtain findings suitable for statistical analysis.

CLINICAL STUDY

Presented at the European Burn Association (EBA), Budapest, Hungary, 2007
and the International Society for Burn Injuries (ISBI), Montreal, Canada, 2008
Burn Centre Rotterdam, the Netherlands

"Itching, is there a remedy ?" a pilotstudy with a rehydration gel-creme in burnpatients

H.J. van Kempen¹ RN, BHSc, J.Dokter² MD, I.M.M.H. Oen³ MD

Burn Centre:nurse and skin therapist(1), medical coordinator(2), research coordinator(3)

Medical Centre Rijnmond-Zuid Rotterdam, the Netherlands; kempenh@mcrz.nl





Introduction

Itching after burns is a matter of body and mind, but little is known regarding its exact mechanism. Results of studies show a gradual decrease in itch intensity in time, but still moderate to serious itch several years after burns (van Loey e.a.). In 2006 a gel-creme that supposed to decrease itch was introduced in our Burn Centre. The main ingredients of Alhydran® are Aloë Vera (Aruba Aloe Barbadenis), vitamins C and E and Jojoba oil. To determine its efficacy we performed a pilotstudy.



Methods

Since November 2006 in the outpatient clinic of the Burn Centre Medical Centre Rijnmond-Zuid itching is scored in patients using the Visual Analogue Thermometer (VAT) or the Numeric Rating Scale (NRS 0-10). Measurements were done before application of Alhydran® and after 1 week, 2 weeks, 1 month and 3 months after the start of treatment. Follow up data were collected on outpatient visit or by phone.



Results

Data analyses with ANOVA and T-test show before starting Alhydran® that the mean VAT/NRS for itch intensity is 6.7(see graphic). There is a decrease in time: the mean score after one week is 3.5, after two weeks 3.6, after one month 2.5 and after three months 2.4. There is no relation between VAT/NRS and gender, age, TBSA%, location, cause of burn or combination therapy with other scar treatment. Besides a decreased VAT score parents of young children mentioned their child was more at ease and had a better night's rest. One patient had discontinued Alhydran® because there was no itch anymore after 1 week and 1 patient reduced antihistaminic medication. Two patients said they stopped the creme because of irritation of the skin.

Conclusions

"Itching can make you crazy". Our first experience indicates that this gel-creme could decrease itch intensity after burns, which contributes to improvement of quality of life. Ongoing research will include additional assessment by a Burns Itch Questionnaire, scar evaluation like the Vancouver Scar Scale, Patients and Observer Scar Assessment Scale, cuto – and dermaspectometrie and a Quality of Life questionnaire.

Reference (1) NEE van Loey, e.a. Itching following burns; epidemiology and predictors; accepted in Burns 2007.

CLINICAL STUDY

Presented at the European Burn Association (EBA), Budapest, Hungary, 2007
Burn Centre, University Hospital Ghent, Belgium

ALHYDRAN

TWO YEARS OF EXPERIENCE WITH A NEW PRODUCT BASED ON ALOE VERA FOR HYDRATION OF BURN SCARS



AUTHORS: S. Monstrey, MD, PhD, A. Pirayesh, MD, E. Lambrecht, RN
S. Lauwaert, RN, J. Verbelen, RN, MN, H. Hoeksema, RC

INSTITUTE: Department of Plastic & Reconstructive Surgery - Burn Centre,
University Hospital Ghent, De Pintelaan 185, 9000 Ghent, Belgium



Introduction: In humans deep dermal burns unfortunately do not heal by regeneration of the damaged skin but by the formation of scar tissue (Fig.1). While the prevention of hypertrophic scars (Fig. 2) with pressure garments and silicone inlays has been widely studied, very little evidence has been published on the optimal skin care after burns. The different structure of scar tissue with the absence of sweat and sebaceous glands results in drier, less elastic and itchy skin. Treatment of these annoying skin problems is of utmost importance for the burn patient. An inquiry in the major European burn centres revealed a plethora of creams and ointments that are used for scar treatment after burns but without any standard therapy. Aruba Aloe Balm (Formula F-BC-096) containing 45% of pure, freshly processed Aruba Aloe Vera gel (Fig. 4) obtained directly from the plant (Fig. 3) was used during the last two years for about 50% of our burn patients. We evaluated the presence of residual defects, elasticity of the skin, subjective feeling of the patient especially itching and ease of use, and the final aesthetic outcome.

Methods: All treated patients children and adults had skin grafts or burns with a conservative healing time of > 18 days. Patients underwent a treatment regimen with pressure and silicone garments. Patients were asked to moisten the scars with a thin layer of the product 3 times a day. In order to avoid biased results the Balm (Formula F-BC-096) was delivered in blanc white tubes (Fig. 8) and patients were not aware of the composition of the product. Follow up visits including clinical assessment and digital photography of the scar were done on a regular basis for at least one year as usual in our centre. For some patients in which a control product for comparison was used measurements for skin elasticity (Dermalab) (Fig. 6) and colour (Dermaspectrometer) (Fig. 5) were performed.



Case 1 : Scar formation after surgery

Start with Alhydran®



3 months with Alhydran®



5 months with Alhydran®



Case 2 : Scald

1 day post burn



Start with Alhydran®



1 year with Alhydran®



Results: No allergic reactions related to the product were noticed. In general the aesthetic outcome of patients treated with the Balm (Formula F-BC-096) was better than in patients treated with other moisturizers as scored in the Vancouver Scar Scale. However these are subjective findings, they were confirmed by objective measurements of the Dermalab and Dermaspectrometer in patients where additionally a control product was used. All patients preferred the test product above other products we always used before in our centre. In more than 90% of the patients itching was reduced as they needed less medication and were more comfortable during the day and especially children had a more quiet sleep. Combination possibilities with pressure garments and silicone sheets are outstanding.

Conclusion: After a survey throughout the burn centres of Europe we noticed that there is definitely no standard for hydration of scars and the use of these moisturizers. After two years of excellent results using this new product we think now there is the possibility to set this standard. Till 2007 this product was not distributed outside the Caribbean. In an attempt to make this product available for all patients in Europe a new brand name **Alhydran®** (Fig.7) was chosen for distribution. A prospective, randomized, double blinded, controlled clinical trial of the **Aruba Aloe Balm (Formula F-BC-096)** in the treatment of split thickness donor sites after wound closure is already registered at www.ClinicalTrials.gov and is recruiting patients.



CASE STUDIES (INDEX)

SURGERY

Surgical intervention	Case study	No. AL20-015-EN	30
-----------------------	------------	-----------------	----

SCARS AND BURNS

Hot water burn	Case study	No. AL20-001-EN	31
Flame burn	Case study	No. AL20-002-EN	32
Second degree burn	Case study	No. AL20-005-EN	33
Scald	Case study	No. AL20-010-EN	34
Chemical explosion	Case study	No. AL20-011-EN	35
Hot water burn	Case study	No. AL20-017-EN	36

AESTHETIC INTERVENTION

Soft surgery	Case study	No. AL20-016-EN	37
--------------	------------	-----------------	----

DONOR SITE

Treatment of donor site	Case study	No. AL20-012-EN	38
-------------------------	------------	-----------------	----

RADIOTHERAPY

Radiotherapy (after care)	Case study	No. AL20-006-EN	39
Treatment during and after radiotherapy	Case study	No. AL20-007-EN	40

CHRONICAL WOUND

Wound edges and peri wound skin	Case study	No. AL20-004-EN	41
---------------------------------	------------	-----------------	-----------

ACTINIC KERATOSIS

Multiple actinic keratosis	Case study	No. AL20-009-EN	42
Erosive skin	Case study	No. AL20-008-EN	43
Actinic Keratosis (after care)	Case study	No. AL20-013-EN	44

CHRONIC SKIN DISORDER

Eczema	Case study	No. AL20-014-EN	45
--------	------------	-----------------	-----------

CASE STUDY

Surgery - Surgical Intervention

Resection of abscess

AL20-015-EN

CASE ALHYDRAN and BAPSCARCARE: Surgical Intervention

1. Case study description

patient age	63 years
gender	female
cause	resection of abscess on palmar side of the wrist
therapy after care	start 12 days after surgical intervention during the day; ALHYDRAN 3 times a day, a very thin layer during the night; BAPSCARCARE S

2. Evolution of the wound



3. Conclusions medical team

- Removal of sutures after 12 days and immediately start of treatment
- The combination of a thin silicone sheet during the night and ALHYDRAN during the day resulted in complete scar maturation within 9 months

Case study: H. Hoeksema, Department of Plastic and Reconstructive Surgery, University Hospital of Ghent, Belgium

CASE STUDY

Scars and burns - Hot water burn

Scald (child)

ALHYDRAN

AL20-001-EN

Case ALHYDRAN: Hot water burn (scald)

1. Case study description

patient age	4,5 years
gender	female
cause	hot water burn
treatment to wound closure	modern wound care dressings
therapy after care	1,5 years after wound closure start with ALHYDRAN, after one month completed with pressure therapy

2. Evolution of the wound



3. Conclusions medical team

- The good hydration by ALHYDRAN improves the flexibility of the vulnerable skin and has a positive effect on itching and redness
- The use of pressure garments (after one month ALHYDRAN therapy) has a good effect on the aspect (flattening) of the scar
- The use of ALHYDRAN contributes significantly to the reduction of itching and redness.
- The scar becomes more flexible

4. Patient experience (and parents)

- Improved flexibility of the skin, less itching and redness
- Clear improvement of the mobility and a visual improvement of the look of the scars

Case study: C. Sukan, Lotus MDC, Ankara, Turkey

CASE STUDY

Scars and burns - Flame burn

Flame burn lower leg

AL20-002-EN

Case ALHYDRAN: Flame burn (lower) leg

1. Case study description

patient age	15 years
gender	female
cause	flame burn
treatment to wound closure	1 week Flammazine, thereafter hydrocolloid gel + Jelonet as wound dressing and light therapy to enhance wound healing
therapy after care	ALHYDRAN, immediately after wound closure pressure garments and Silicone sleeve + hydration with ALHYDRAN (2 to 3 times a day) after one year follow up with ALHYDRAN alone

2. Evolution of the wound



3. Conclusions medical team

- Wound healing after 19 days
- Evolution to complete maturation within 1 year, with excellent results regarding to color (slight hypo pigmentation) and elasticity of the scar
- Absolutely no hypertrophic scarring and impairment of mobility

4. Patient experience (and parents)

- The patient was originally not very excited by the typical Aloe Vera smell of the ALHYDRAN
- However this patient was very satisfied by the other properties of the cream such as easiness of application and the hydration after application, resulting in much better suppleness of the skin
- Less itching was noticed by this young girl

Case study: Prof. S. Monstrey and H. Hoeksema, University Hospital Ghent – department of Plastic and Reconstructive Surgery and Burns Center, Belgium

CASE STUDY

Scars and burns - Second degree burn

Firework burn

ALHYDRAN

AL20-005-EN

Case ALHYDRAN: Second degree burn

1. Case study description

patient age	45 years
gender	male
cause	fireworks burn
treatment to wound closure	Flammazine
therapy after care	5 weeks post accident start with ALHYDRAN, 3 daily doses, brought on thick during the first few weeks

2. Evolution of the wound



3. Conclusions medical team

- Deep second degree burn with serious ankle damage
- Therapy starts with Flammazine, patient evolves well, only a lot of complaints on itching of closed skin
- After 5 weeks: start with ALHYDRAN on the freshly closed skin, 3 times per day
- ALHYDRAN had good result on the skin, which improved rapidly
- Itching complaints disappeared immediately when ALHYDRAN was used

4. Patient experience

- Flammazine gave a lot of itching problems
- Upon start of the treatment with ALHYDRAN, immediate reduction of itching
- Therapy: 3 daily doses, or each time the skin was itching
- After 6 months, the skin was soft and calm, patient started sporting again
- Skin looks good and calm now
- After 18 months, maintenance of 1 daily dose of ALHYDRAN

Case study: K. Veel MD PhD, AAB Aruba

CASE STUDY

Scars and burns - Scald

Hot water burn from exploding radiator

AL20-010-EN

Case ALHYDRAN: Scald (hot water from exploding radiator)

1. Case study description

patient age	52 years
gender	male
cause	scald (hand, face, chest)
treatment	ALHYDRAN (twice a day)

2. Evolution of the skin



Day 1 after scald. Blisters punctured / commencing with ALHYDRAN as single therapy



Day 7 following treatment ALHYDRAN



Day 14 following treatment ALHYDRAN

3. Conclusions medical team

- Positively surprised at fast results & healing process
- Patient indicated absence of pain and itching
- Uneventful healing process within 4 weeks

4. Patient experience

- No pain or itching
- Rapid healing, can resume activities
- Easy application ALHYDRAN, no fragrance

Case study: Dr R. Ceulen, Dermatologist, Albert Schweitzer Hospital, Dordrecht, the Netherlands

CASE STUDY

Scars and burns - Chemical explosion
Second to third-degree burns



AL20-011-EN

Case ALHYDRAN: Burn caused by chemical explosion

1. Case study description

patient age	58 years
gender	male
cause	burn scar after explosion
treatment to wound closure	Flamazine and daily dressing change
therapy after care	ALHYDRAN, pressure gloves, scar therapy by skin therapist, endermotherapy

2. Evolution of the skin



Start ALHYDRAN



After 3 months of ALHYDRAN therapy



After 6 months of ALHYDRAN therapy

3. Conclusions medical team

- Fast healing
- Improved restoration skin hydration
- Visibly improved supple skin
- Reduction squamae, ie. skin particles

4. Patient experience

- Skin has stopped itching
- Improved flexibility of the skin
- Reduced contractions
- Easy to apply

Case study: Ms J.M. de Graaf and Ms C.T.M. Rijkenberg, Practice Skin Therapists, Oosterbeek, the Netherlands

CASE STUDY

Scars and burns - Hot water burn

Deep second degree burns

AL20-017

Case ALHYDRAN and SCARBAN: Hot water burn

1. Case study description

patient age	84 years
gender	female
cause	Trauma after stumbling with a pot of hot tea. The patient landed on her knees and the pot of hot tea fell over the patient's left shoulder.
why scar treatment	Development of a hypertrophic scar after deep second degree burns on the patient's back. This scar caused itching and pain with moving.
start scar treatment	3 and a half months after trauma
therapy	<ul style="list-style-type: none">- ALHYDRAN and SCARBAN for a duration of 13 months.- SCARBAN Elastic sheets were used for 12 hours during the day.- In the evening the sheets were removed. At that time ALHYDRAN was applied for the night.- In the morning after showering the SCARBAN Elastic sheets were re-applied.

2. Evolution of the scar



3. Conclusions medical team

- Results above expectations
- Nice improvement of the quality of life for the patient.
- Scar quality has clearly improved and functionality improved.

4. Patient experience

- Patient is very satisfied after treatment
- Changes of the silicon sheets were time-consuming at times, but they knew what they did it for.
- Patient does not experience any itching or pain anymore.

Author: E. Harink, MSc, Nurse Practitioner, Department Vascular Surgery, Hospital Isala Zwolle, a Dutch Case Report

CASE STUDY

Aesthetic intervention - Soft surgery

PlexR treatment

ALHYDRAN

AL20-016-EN

Case ALHYDRAN: Soft surgery aftercare

1. Case study description

patient age	54 year
gender	female
cause	soft surgery to firm up skin and reduce wrinkles
treatment	PlexR treatment
aftercare therapy	- start applying ALHYDRAN immediately after PlexR treatment (2 to 3 times daily) - light cooling of the skin during the first 4 days (do not wet) - dab on after 4 days (do not rub)

2. Evolution of the skin



Before treatment



Start ALHYDRAN immediately after treatment



2 days after treatment: scab development



6 days after treatment: reddened skin without scabs



1 month after treatment: light reddening



5 months after treatment: fully recovered

3. Conclusions medical team

We perform a number of aesthetic interventions in our clinic using a PlexR device. High-energy plasma is used to burn the skin, resulting in skin contraction. Alhydran® is used for aftercare.

- Wound healing is accelerated. The period of skin redness after the treatment is significantly reduced
- Much less hyperpigmentation
- The healing process is more efficient than with aqueous products. An aqueous cream causes scabs to fall off too quickly.
- The healed wound is virtually scar-free
- Strong reduction of itching complaints following the treatment

4. Patient experience

- Alhydran is easy to apply (spreads well and is quickly absorbed)
- Recovery feels and is visibly rapid
- Flexible skin after treatment

Case study: MD E.R. Berends, Perfect Look Clinic, Westerbork, the Netherlands

CASE STUDY

Donor site - Treatment of donor site

AL20-012-EN

Case ALHYDRAN: Donor site

1. Case study description

patient age	13 years
gender	female
cause	donor site
treatment to wound closure	epidermal skin substitute
therapy after care	ALHYDRAN, immediately after wound closure pressure garments and hydration with ALHYDRAN (2 to 3 times a day)

2. Evolution of the wound



3. Conclusions medical team

- Full wound closure within 14 days
- Evolution to complete maturation within 1 year, with excellent results regarding to color (slight hypo pigmentation) and elasticity of the donor site
- Significantly less redness and itching during maturation

4. Patient experience (and parents)

- No significant pain during wound healing
- No significant itching during the treatment, before nor after wound closure
- Good flexibility and feel of the skin on the donor site

Case study: Prof. S. Monstrey and H. Hoeksema. University Hospital of Ghent – department of Plastic and Reconstructive Surgery and Burns Centre, Belgium

CASE STUDY

Radiotherapy - Radiotherapy (after care)

Mastectomy with axillary dissection



AL20-006-EN

Case ALHYDRAN: Radiotherapy (after care)

1. Case study description

patient age	68 years
gender	female
diagnosis	mastectomy with axillary dissection
therapy after care	started with ALHYDRAN (3 times daily) immediately after radiotherapy

2. Evolution of the skin



3. Conclusions medical team

- ALHYDRAN ensures that the radiated area is less inflamed, red and painful
- ALHYDRAN ensures that the skin heals optimally. As a result of the good and rapid healing, the follow-up phase can be started earlier
- ALHYDRAN is pleasant to use and simple to apply

4. Patient experience

- ALHYDRAN ensures less pain and redness
- There is noticeably rapid improvement to the radiated area
- ALHYDRAN is pleasant to use

Case study: Ms M. voor 't Hekke, Skin Therapist, Zwolle, the Netherlands

CASE STUDY

Radiotherapy - Treatment during and after radiotherapy
Breast carcinoma


AL20-007-EN

Case ALHYDRAN: Radiotherapy

1. Case study description

patient age	42 years
gender	female
diagnosis	breast carcinoma
therapy after care	ALHYDRAN during and after the radiation period

2. Evolution of the skin



At the end of radiation



2 months after radiation and treatment with ALHYDRAN

3. Conclusions medical team

- ALHYDRAN is a good alternative for cetomacrogol
- Patients are almost always positive about its use
- ALHYDRAN is better than an Aloe Vera gel only, due to hydrating properties of ALHYDRAN
If only a gel is used, the radiated skin will become too dry and flaky
- ALHYDRAN is used with grade I and II radiation dermatitis

4. Patient experience

- ALHYDRAN is better than an indifferent cream (cetomacrogol), because ALHYDRAN eliminates the pain of radiation skin burn
- ALHYDRAN reduces itching
- ALHYDRAN was easy to apply and has a neutral odour

Case study: N.S. Russell, radiotherapist, Netherlands Cancer Institute (NKI), Antoni van Leeuwenhoek Hospital, Amsterdam, the Netherlands

CASE STUDY

Chronical wound - Wound edges and peri wound skin

Chronical wound after open fracture luxation



AL20-004-EN

Case ALHYDRAN: Chronical wound

1. Case study description

patient age	79 years
gender	female
cause	open fracture luxation, right ankle with osteosynthesis material post operation, a chronical wound developed
treatment to wound closure	- curettage and hydrogel on wound - wound borders with ALHYDRAN
therapy after care	ALHYDRAN

2. Evolution of the wound



3. Conclusions medical team

- After luxation fracture and the operation with osteosynthesis material, the wound had never been closed
- Due to the presence of the characteristics of a chronical wound with underlying OS material, there is a big chance of deeper infections and the wound will be very difficult to close
- OS material was removed laterally
- The wound borders have improved significantly due to treatment with ALHYDRAN
- In the after care only ALHYDRAN was used

Case study: E. Roovers, Wound Care Coordinator, Ziekenhuisnetwerk Antwerpen (ZNA), Belgium

CASE STUDY

Actinic keratosis - Multiple actinic keratosis

Serious skin reaction after 5-ala PDT treatment


AL20-009-EN

Case ALHYDRAN: Multiple actinic keratosis

1. Case study description

patient age	81 years
gender	male
diagnosis	- actinic keratosis of the scalp - day 1 after PDT, patient was seen during an emergency consultation with serious reactions: inflamed erythema, erosions, crusting
treatment	a single session of 5-ala PDT
therapy after care	ALHYDRAN twice daily

2. Evolution of the skin



Start of ALHYDRAN as a single treatment, applied twice a day



Day 7 after starting ALHYDRAN

3. Conclusions medical team

- Treatment with ALHYDRAN gives a good and quick clinical improvement (within a week)

4. Patient experience

- Complaints of pain disappeared within 48 hours
- ALHYDRAN had a cooling effect

Case study: Dr R. Ceulen, Dermatologist, Albert Schweizer Hospital, Dordrecht, the Netherlands

CASE STUDY

Actinic keratosis - Erosive skin

Erosive skin after 5-Fluorouracil treatment



AL20-008-EN

Case ALHYDRAN: Erosive skin

1. Case study description

patient age	70 years
gender	male
diagnosis	actinic keratosis (skin cancer)
treatment	3 weeks twice daily application of 5-Fluorouracil
therapy after care	after 3 weeks start ALHYDRAN

2. Evolution of the skin



3. Conclusions medical team

- ALHYDRAN is easy to apply to mildly erosive skin
- ALHYDRAN is a pleasant alternative to fusidic acid cream
- ALHYDRAN does not entail a risk of bacterial resistance, unlike fusidic acid cream
- ALHYDRAN keeps the skin slightly oily and hydrated
- ALHYDRAN also promotes wound healing

4. Patient experience

- ALHYDRAN is easy to apply
- Nice to use, does not sting
- Noticeable result in just a couple of days
- ALHYDRAN also works well on grazes

Case study: Dr D. Njoo, dermatologist, Hospital Group Twente, Hengelo (Ov), the Netherlands

CASE STUDY

Actinic keratosis - Treatment during and after radiotherapy

Treated with cryotherapy

AL20-013-EN

Case ALHYDRAN: Actinic keratosis

1. Case study description

patient age	72 years
gender	male
diagnosis	actinic keratosis
treatment	patient had 2 actinic keratosis on the scalp both lesions were treated with 2 cycles of liquid nitrogen spray treatment
therapy after care	ALHYDRAN cream twice daily for 14 days

2. Evolution of the wound



Start ALHYDRAN treatment, 2x daily



14 days after start of ALHYDRAN treatment, 2x daily

3. Conclusions medical team

- The treated skin areas closed after 14 days, leaving no scar formation or hypopigmentation
- ALHYDRAN is easy to use and has no local side effects
- ALHYDRAN can be used as an alternative for fusidinic acid cream
- In contrast to fusidinic acid, ALHYDRAN does not cause bacterial resistance
- ALHYDRAN cream proved to heal wounds rapidly

4. Patient experience

- Easy in use
- No skin irritation or other side effects
- Fast clinical results, within a few days

Case study: Dr D. Njoo, dermatologist, Hospital Group Twente, Hengelo (Ov.), the Netherlands

CASE STUDY

Chronic skin disorder - Eczema

Atopic eczema

AL20-014-EN

Case ALHYDRAN: Atopic eczema

1. Case study description

patient age	22 years
gender	female
diagnosis	atopic eczema patient had suffered from atopic eczema since childhood, the patient was reluctant to use local corticosteroids as she feared atrophy of the skin. The eczema is quite persistent and recurring, especially inside the elbows
treatment	ALHYDRAN cream twice daily, monotherapy inside the elbows

2. Evolution of the skin



3. Conclusions medical team

- The eczema decreased strongly within 2 weeks
- Monotherapy with ALHYDRAN cream proven to be effective in mild eczema cases
- ALHYDRAN delivered results comparable with a local corticosteroid
- ALHYDRAN had no side effects, in particular no skin atrophy

4. Patient experience

- Easy to apply and quickly absorbed
- No skin irritation or other side effects
- More confidence in the product, as there are no side effects such as atrophy

Case study: Dr Njoo, dermatologist, Hospital Group Twente, Hengelo (Ov.), the Netherlands

LITERATURE REFERENCES

	Author(s)	Title	Publication
1	Hoeksema H, de Vos M, Verbelen J, Pirayesh A, Monstrey S.	Scar management by means of occlusion and hydration: A comparative study of silicones versus a hydration gel-cream.	Burns 2013
2	Klotz T, Kurmis R, Munn Z, Heath K, Greenwood J.	Moisturisers in scar management following burn: A servery report.	Burns 2017
3	Rondas AALM, Schols JMGA	Aplication of medical moisture retention cream (Alhydran), a new option in the treatment of venous eczema.	J Gerontol Geriatr Res 2017
4		Available case studies (see pages 28 - 45)	
5	Pirayesh A, Hoeksema H, Richters C, Verbelen C, Monstrey S.	Glyaderm dermal substitute: Clinical application and long-term results in 55 patients.	Burns 2014
6	Monstrey S, Pirayesh A, Lambrecht E, Lauwaert S, Verbelen J, Hoeksema H.	Two years of experience with a new product based on aloe vera for hydration of burn scars.	EBA 2007
7	Hoeksema H, Van Gastel K, Van Geertruy V, Lafaire P, Verbelen J, Monstrey S.	Development of a new brochure making aftercare of burns more understandable for patients.	EBA 2013
8	Middelkoop E, Monstrey S, Téot L, Vranckx JJ.	Scar Management Practical Guidelines. Maca-Cloetens: 2011;1-109.	2011
9	Mustoe TA.	Review - Evolution of Silicone Therapy and Mechanism of Action in Scar Management. 32:82-92: fig. 2- page 89, page 88.	Aesth Plast Surg 2008
10	Njoo MD.	Management of non-melanoma skin cancer, care and after care. ISBN 978-90-821921-0-0.	Book publication 2014
11	Ziegenthaler H.	Effect of scar treatment products on unpleasant physical manifestations in patients suffering from extensive burn scarring.	DAV 2010
12	Ziegenthaler H.	Scars after large surface thermal burns – registration of quality of life and therapeutic influence of ALHYDRAN.	EBA 2011
13	Giovannini U.	Application of ALHYDRAN in the treatment of pathologic scars after cryoshape therapy.	EBA 2011
14	Ashman S, Kray SJ, Tyne & Wear.	Psychological wellbeing – scar.	EWMA 2014
15	van Kempen HJ, Dokter J, Oen IMM.	Itching, is there a remedy? A pilotstudy with a rehydration gel-crème in burnpatients.	EBA 2007 ISBI 2008
16	Rijkenberg R.	Use of ALHYDRAN cream in the treatment of burn scars following a chemical explosion – a case study.	EBA 2015
17	Rondas AALM, Vestjens JA, Schols JMGA.	ALHYDRAN, a new option in the treatment of hypostatic eczema	WUWHS 2017
18	Moorgat P, Anthonissen M, Meirte J, Van Daele U, Vanhullebusch T, Lafaire C, De Cuyper L, Maertens K.	The effect of a topical cream application on water distribution in healthy skin and burn scars: pilot project	EBA 2018
19	Monstrey S, Middelkoop E, Vranckx JJ, Bassetto F, Ziegler UE, Meaume S, Téot L.	Updated scar management practical guidelines 2014 67, 1017-1025	J of Plastic Reconstructive & Aesthetic Surgery 2014
20	Van Leen MWF, Hoeksema H, Schols JMGA	MMRC (Alhydran), an interesting treatment option for a disrupted skin barrier	J Gerontol and Geriatr Medicine 2021
21	Hofland H.W.C.	The occurrence of itch after burns: is there a need for oral medication?	EBA 2023